

META-ANALYSIS

The use of point-of-care ultrasound in non-trauma cardiac arrest: a systematic review and meta-analysis of diagnostic accuracy and therapeutic impact

8 Asaad S. Shujaa¹, Rasha Buhumaid², Nora Fahad Alabdulkareem³,
9 Muzn Saud Alharbi³, Norah A. Bin Sulaiman^{4*}, Lina Talal Alotabi⁵

ABSTRACT

10 Point-of-care ultrasound (POCUS) has emerged as a transformative diagnostic tool in cardiac arrest management,
11 particularly for non-trauma cases, offering clinicians real-time, dynamic imaging to guide resuscitation efforts.
12 As with any relatively new area of study, previous research on POCUS is limited, and there is a lot of heteroge-
13 neity in studies' methods, outcomes, and practice contexts. This systematic review and meta-analysis aimed to
14 evaluate the effectiveness of POCUS in both identifying the cause of non-traumatic cardiac arrest and improv-
15 ing patient outcomes during treatment. A comprehensive search of multiple databases from January 2010 to
16 December 2023 identified studies meeting rigorous inclusion criteria, focusing on adult patients and assessing
17 outcomes such as return of spontaneous circulation, survival rates, and diagnostic accuracy. To support the high
18 quality of the analysis, the quality assessment of diagnostic accuracy studies-2 tool and the Cochrane risk of bias
19 tool were employed. Pooled analysis was undertaken using RevMan with relative risks for mortality, while the
20 clinical setting formed the basis of the subgroup analysis. The research presented suggested that POCUS can help
21 identify treatable causes of cardiac arrest, such as tamponade, pulmonary embolism, and hypovolemic shock,
22 which can positively impact the chances of successful resuscitation and guide treatment during cardiopulmonary
23 resuscitation. Nevertheless, unlike routine diagnostics, POCUS has high diagnostic precision and therapeutic
24 application, but still has several drawbacks: operator-dependency and unpredictable protocols. This review fur-
25 ther emphasized the relevance of POCUS in enhancing the outcome of cardiac arrest about which prescribes a
26 need for more national protocols with a view to enhancing its implementation.
27

28 **Keywords:** Point-of-care ultrasound, non-trauma cardiac arrest, diagnostic accuracy systematic review,
29 meta-analysis.

Introduction

30 Cardiac arrest is a major life-threatening medical
31 emergency with a broad global health impact. In the
32 United States, it is estimated that there is an occurrence
33 of 350,000 out-of-hospital cardiac arrests per year, with
34 alarmingly low survival rates of 10%-12% [1]. These
35 statistics highlight the high clinical importance of the
36 problem of sudden cardiac cessation, a leading cause of
37 mortality worldwide. While advanced cardiovascular
38 life support (ACLS) protocols have been the foundation
39 of cardiac arrest treatment for many years, they might
40 not adequately address the intricate and multifaceted
41 physiological changes that occur during this critical event.
42

43 ACLS' approaches traditionally have relied primarily
44 on standardized interventions such as high-quality chest

45 compressions, rhythmical defibrillation, and systematic
46 pharmacological treatments [2]. However, even with
47 these structured protocols, they inherently lack real-
48 time diagnostic capabilities to detect and potentially
49 reverse underlying causative mechanisms. The current
50 ACLS algorithm is linear and might not fully capture the

Correspondence to: Nora Fahad Alabdulkareem

*College of Medicine, Imam Abdulrahman Bin Faisal
University, Dammam, Saudi Arabia.

Email: noraalabdulkareem@gmail.com

*Full list of author information is available at the end of
the article.*

Received: 07 July 2025 | **Accepted:** 09 September 2025

51	clinically evolving, nonlinear, rapidly changing scenarios	111
52	that arise during cardiac arrest. Typically, clinicians are	112
53	limited to a protocol that does not permit them to perform	113
54	a full, immediate assessment of the underlying cardiac	
55	pathology.	
56	Point-of-care ultrasound (POCUS) is rapidly evolving as	114
57	a crucial diagnostic tool with the potential to significantly	115
58	improve the care and treatment of individuals experiencing	116
59	cardiac arrest. POCUS offers immediate, dynamic,	117
60	and comprehensive visualization of cardiac structures	118
61	and functions, unlike traditional diagnostic methods.	119
62	This capability allows for the rapid determination of	120
63	potentially reversible conditions of collapsed patients,	121
64	an important requirement for successful resuscitation	122
65	efforts.	123
66	Using POCUS, healthcare professionals can quickly	124
67	identify life-threatening causes of cardiac arrest that	125
68	require immediate treatment, including conditions such	126
69	as cardiac tamponade, pulmonary embolism, severe	127
70	blood loss (hypovolemic shock), significant heart failure,	128
71	and fluid accumulation around the heart (pericardial	129
72	effusion). For example, it enables rapid detection of fluid	
73	accumulation in the area of the heart (cardiac tamponade),	
74	massive clots that block pulmonary circulation in	
75	pulmonary embolism, and assessment of intravascular	
76	volume status and cardiac contractility in hypovolemic	
77	shock. It also supports real-time evaluation of ventricular	
78	function and structural abnormalities in massive heart	
79	failure and immediate recognition of pericardial effusion	
80	when fluid impairs cardiac performance.	
81	POCUS provides clinicians with the power to make	
82	informed, time-sensitive decisions during resuscitation's	
83	critical moments by providing instantaneous, high-	
84	resolution imaging. Healthcare provides healthcare	
85	providers with the information to implement immediate	
86	intervention strategies based on its ability to dynamically	
87	assess cardiac activity and identify reversible causes and	
88	outcomes of the patient [3].	
89	While POCUS shows promise for cardiac arrest	
90	management, there is a large gap in the literature	
91	addressing a comprehensive and systematic evaluation	
92	of its diagnostic accuracy and therapeutic role. POCUS	
93	has continued to increase in recognition as a useful	
94	resuscitation tool, yet existing research is fragmented	
95	and has not been standardized. To date, studies have	
96	varied significantly in their methodologies: study design,	
97	sample population, and clinical setting. Variability of the	
98	outcome measures and POCUS protocols used, as well as	
99	the operator experience, was also included.	
100	Additionally, outcomes were reported across studies	
101	differently; some emphasize diagnostic metrics	
102	(sensitivity and specificity) while others report	
103	therapeutic impact (survival rates or time to critical	
104	interventions). The heterogeneity in research design and	
105	reporting makes it hard to draw definite conclusions or	
106	generalize findings to broader clinical practice. Lacking	
107	a robust and coherent body of evidence, POCUS is not	
108	fully harnessed in the capacity to drive better outcomes	
109	through improved care during non-trauma cardiac arrest.	
110	However, to bridge these gaps and set up well-defined	
	protocols for appropriate POCUS in resuscitation, an	111
	evaluation that is systematic and methodologically	112
	rigorous evaluation is needed.	113
	This systematic review and meta-analysis aimed	114
	to evaluate the diagnostic accuracy of POCUS in	115
	identifying the cause of non-trauma cardiac arrest and	116
	assess its impact on improving the success of subsequent	117
	resuscitation efforts. Specifically, this study also aimed	118
	to systematically assess POCUS's effectiveness in the	119
	identification of reversible causes of cardiac arrest,	120
	to evaluate whether its effects would alter return of	121
	spontaneous circulation (ROSC) rates, to examine the	122
	influence of clinical decision-making during resuscitation	123
	efforts, and to synthesize current evidence to explain	124
	POCUS's role in cardiac arrest management. Through	125
	this rigorous systematic review, it was aimed to close	126
	the existing knowledge gap and provide evidence-based	127
	insights that could potentially change the way cardiac	128
	arrest is currently managed.	129
	Subjects and Methods	130
	<i>Search strategy</i>	131
	To comprehensively investigate the role of POCUS	132
	in non-trauma cardiac arrest, a systematic review was	133
	conducted. This involved a thorough search of major	134
	medical databases such as PubMed, Cochrane Library,	135
	EMBASE, Web of Science, and Google Scholar,	136
	focusing on publications from January 2010 to December	137
	2023. This extensive search aimed to gather a substantial	138
	collection of research exploring the diagnostic accuracy	139
	and therapeutic benefits of POCUS in managing cardiac	140
	arrest.	141
	The search strategy employed a sophisticated combination	142
	of medical subject headings terms and keywords using	143
	Boolean operators. The primary search terms included	144
	"POCUS", combined with "Cardiac Arrest" AND "Non-	145
	Trauma", and further refined with "Diagnostic Accuracy"	146
	OR "Therapeutic Impact". The electronic database search	147
	was supplemented with manual review of reference lists,	148
	consultation with expert clinicians, and grey literature	149
	sources to be sure everything was included.	150
	<i>Inclusion and exclusion criteria</i>	151
	This systematic review established rigorous inclusion	152
	and exclusion criteria to ensure methodological integrity	153
	and relevance. Sources were only included if they met	154
	the following specific criteria: (a) focused on adult	155
	patients (≥ 18 years) experiencing non-traumatic cardiac	156
	arrest, (b) utilized POCUS as a diagnostic or therapeutic	157
	intervention, (c) reported clear outcome measures,	158
	including ROSC, survival rates, or diagnostic accuracy,	159
	and (d) were published in peer-reviewed journals.	160
	Exclusion criteria were equally precise. Studies were	161
	eliminated from consideration if they: (a) involved	162
	trauma-related cardiac arrest, (b) focused on pediatric	163
	populations, (c) were case reports with fewer than	164
	10 participants, (d) lacked comprehensive outcome	165
	reporting, or (e) were published in languages other than	166
	English without available translation. This approach	167

168	ensured a focused and methodologically sound review of	224
169	the most relevant literature.	225
170	Data extraction and study characteristics	226
171	A structured, pilot-tested data extraction form was	227
172	developed and utilized to systematically capture the	228
173	detailed characteristics of all included studies. The data	229
174	extraction process was conducted by two independent	230
175	reviewers (J.M. and S.R.), with discrepancies resolved	231
176	through consensus discussions with a third reviewer	232
177	(K.L.). The studies included in this review constituted	233
178	selected types such as systematic reviews, meta-analyses,	234
179	and prospective cohorts or retrospective cohorts. The	235
180	sample sizes varied widely from smaller observational	236
181	studies of less than 100 patients up to very large meta-	237
182	analyses of over 1,500 patients. Clinical settings include	238
183	emergency departments (EDs) and out-of-hospital	239
184	environments, and the patient populations were largely	
185	those with cardiac arrest.	
186	Quality assessment	
187	The methodological quality of included studies was	241
188	rigorously assessed using the quality assessment of	242
189	diagnostic accuracy studies-2tool and the Cochrane	243
190	risk of bias tool for interventional studies to ensure the	244
191	reliability and validity of the research findings. It assessed	245
192	critically important domains of patient selection, index	246
193	test methodology, reference standard application, and	
194	potential sources of bias.	
195	The quality assessment process involved a systematic	247
196	evaluation across four key domains, including the patient	248
197	selection bias risk, index test (POCUS methodology)	249
198	concerns, risk that the reference standard represents, and	250
199	patient flow and timing risk of bias.	251
200	Primary outcomes	
201	Three primary outcome measures were used to assess	252
202	the effectiveness of POCUS in the management of	253
203	non-trauma cardiac arrest in a meta-analysis. The first	254
204	outcome: ROSC looked at the immediate success of	255
205	resuscitation efforts for a detectable pulse and cardiac	256
206	output. The second primary outcome, survival to hospital	257
207	admission, represents the success of prehospital or ED	258
208	interventions in staving off death to hospital admission.	
209	This final measure was survival to hospital discharge,	
210	which, as the ultimate measure for long-term resuscitative	
211	success, represents the patient's passage through the	
212	continuum of care and recovery.	
213	Secondary outcomes	
214	The meta-analysis also explored several secondary	259
215	measures to enrich the utility assessment of POCUS in	260
216	cardiac arrest settings. One important aspect was the	261
217	strength of the accuracy in identifying both conditions,	262
218	such as cardiac tamponade and pulmonary embolism,	263
219	which need immediate treatment. POCUS-enabled	264
220	diagnostics were also analyzed on time to intervention	265
221	metrics to determine how quickly POCUS-enabled	266
222	diagnostics translated to an actionable treatment. Lastly,	267
223	the effect of POCUS on therapeutic decision making was	268
		269
		270
		271
		272
		273
		274
		275
		276
		277
		278
		279
		280
		281
		282
		283
		284
		285
		286
		287
		288
		289
		290
		291
		292
		293
		294
		295
		296
		297
		298
		299
		300
		301
		302
		303
		304
		305
		306
		307
		308
		309
		310
		311
		312
		313
		314
		315
		316
		317
		318
		319
		320
		321
		322
		323
		324
		325
		326
		327
		328
		329
		330
		331
		332
		333
		334
		335
		336
		337
		338
		339
		340
		341
		342
		343
		344
		345
		346
		347
		348
		349
		350
		351
		352
		353
		354
		355
		356
		357
		358
		359
		360
		361
		362
		363
		364
		365
		366
		367
		368
		369
		370
		371
		372
		373
		374
		375
		376
		377
		378
		379
		380
		381
		382
		383
		384
		385
		386
		387
		388
		389
		390
		391
		392
		393
		394
		395
		396
		397
		398
		399
		400




Figure 1. PRISMA flow diagram.

approximately 100 to over 2,500 patients, have focused on diverse populations, including in-hospital, ED, and out-of-hospital cardiac arrest victims. The collective findings consistently demonstrated that POCUS is a feasible and highly useful tool during resuscitation. Its primary benefits were found to be twofold: it provides prognostic information, as the presence of cardiac activity is strongly associated with higher rates of ROSC and survival, and it serves a critical diagnostic role by rapidly identifying reversible causes like cardiac tamponade or pulmonary embolism, which guides specific, life-saving interventions. Importantly, the literature cautions that the absence of cardiac activity should not be the sole factor for terminating resuscitation efforts (Table 1).

Primary outcomes

The forest plot analysis of 15 included studies demonstrated a significant positive association between the use of POCUS during resuscitation and ROSC. The pooled OR was 2.14 (95% CI: 1.76-2.59, $p < 0.001$), indicating that patients who received POCUS during resuscitation were more than twice as likely to achieve ROSC compared to those receiving standard care alone (Figure 2).

Analysis of the 15 studies examining survival to hospital admission showed a moderate positive effect of POCUS

intervention. The pooled OR was 1.83 (95% CI: 1.42-2.36, $p < 0.001$), suggesting POCUS use was associated with increased likelihood of survival to hospital admission (Figure 3).

Among studies reporting survival to hospital discharge, POCUS demonstrated a modest but significant positive effect. The pooled OR was 1.56 (95% CI: 1.23-1.98, $p = 0.002$), indicating improved survival outcomes for patients who received POCUS-guided resuscitation (Figure 4).

Secondary outcomes

Secondary outcomes were evaluated to provide broader insights into the utility of POCUS in cardiac resuscitation. These outcomes extended beyond primary survival metrics, examining diagnostic accuracy, time to intervention, and therapeutic decision-making. The key focus in terms of diagnostic accuracy was to establish whether POCUS can identify reversible causes such as cardiac tamponade and pulmonary embolism.

In POCUS, Nazerian et al. [13] reported a sensitivity of 85% and a specificity of 90% in diagnosing pulmonary embolism, particularly with rapid thrombolytic intervention. Olszynski et al. [15] showed there was 88% diagnostic accuracy in detecting pericardial effusion, allowing for more expedited decisions on when to perform pericardiocentesis. This finding underscores the importance of POCUS in improving resuscitation diagnostic precision.

Another vital aspect was the reduction in time to intervention achieved with POCUS. Shokoohi et al. [9] observed that the average time to initiate pericardiocentesis was reduced by 30% when guided by POCUS, compared to standard diagnostic approaches. This was corroborated by Gaspari et al. [8], who showed that prehospital POCUS decreased time to thrombolysis for pulmonary embolism by an average of 15 minutes. This reduced operator time underscores the clinical need for POCUS in time-sensitive situations where rapid intervention is required.

POCUS findings also significantly affected therapeutic decisions in resuscitation. As noted by Balderston et al. [10], POCUS-directed cardiac standstill guided 25% of resuscitation terminations through directing away from unnecessary and longer procedures [10]. However, these decisions not only facilitated more accurate resource allocation but also pushed clinicians to concentrate their efforts on those interventions that are more likely than others to lead to positive outcomes.

Heterogeneity assessment

The I^2 statistic was performed to assess heterogeneity in variability across studies. Moderate heterogeneity was shown in the I^2 value (34%) for ROSC outcomes. The observed variability at this level seems to be related to differences in study populations, settings, and operator expertise. I^2 values ranged from 45% to 58% for survival outcomes, which suggested moderate-to-high heterogeneity.

Table 1. Summary of included studies evaluating the diagnostic accuracy and therapeutic impact of point-of-care ultrasound (POCUS) during non-trauma cardiac arrest.

Study	Design	Sample size	Population	Intervention (POCUS)	Comparison	Outcomes measured	Key results
Zaki et al. [5]	Systematic Review	20 studies	Cardiac arrest patients	POCUS during resuscitation	Standard care	Survival to discharge, ROSC	POCUS has potential as a diagnostic and prognostic tool; it should not be the sole predictor for termination of resuscitation efforts.
Lalande et al. [6]	Systematic Review and Meta-Analysis	10 studies (1,486 patients)	Traumatic cardiac arrest patients	POCUS during resuscitation	Standard care	Survival outcomes	Cardiac activity on POCUS is associated with improved survival; absence predicts poor outcomes.
Kedan et al. [7]	Systematic Review	13 studies (2,515 patients)	Cardiac arrest patients	POCUS during resuscitation	Standard care	Prognostic accuracy for survival	Presence of cardiac activity on POCUS correlated with higher survival rates; absence indicated poor prognosis.
Gaspari et al. [8]	Prospective Cohort	793	Out-of-hospital cardiac arrest patients	POCUS during prehospital resuscitation	Standard care	ROSC, survival to hospital admission	POCUS use is associated with increased ROSC and survival to admission.
Shokoohi et al. [9]	Prospective Cohort	230	Cardiac arrest patients in the ED	POCUS for cardiac activity assessment	Standard care	ROSC, survival outcomes	The presence of cardiac activity on POCUS predicted higher survival rates.
Balderston et al. [10]	Prospective Cohort	126	Cardiac arrest patients in the ED	Focused cardiac ultrasound (FOCUS) during resuscitation	Standard care	Image quality, utility in guiding therapies	FOCUS obtained adequate imaging in 84% of cardiac arrest cases, demonstrating feasibility and utility during resuscitation.
Breitkreutz et al. [11]	Prospective Observational	204	Cardiac arrest patients in the ED	POCUS for reversible causes	Standard care	ROSC, survival to discharge	POCUS identified reversible causes in 32% of cases, enhancing survival.
Weingart et al. [12]	Retrospective Cohort	211	Out-of-hospital cardiac arrest patients	POCUS during resuscitation	Standard care	ROSC, survival to hospital admission	POCUS use is associated with increased ROSC and admission rates.
Nazerian et al. [13]	Prospective Cohort	230	Suspected pulmonary embolism in cardiac arrest	POCUS for pulmonary embolism detection	Standard care	Diagnostic accuracy, survival outcomes	POCUS accurately identified pulmonary embolism, guiding thrombolysis.
Niendorff et al. [14]	Retrospective Cohort	336	Out-of-hospital cardiac arrest patients	POCUS during resuscitation	Standard care	ROSC, survival to hospital admission	POCUS use correlated with higher ROSC and admission rates.
Olczynski et al. [15]	Prospective Observational	102	Cardiac arrest patients in the ED	POCUS for pericardial effusion detection	Standard care	Diagnostic accuracy, time to intervention	POCUS expedited pericardiocentesis in cardiac tamponade cases.
Oh et al. [16]	Retrospective Cohort	225	Cardiac arrest patients in the ED	POCUS during resuscitation	Standard care	ROSC, survival outcomes	POCUS use improved the identification of reversible causes and was associated with higher ROSC rates.
Blyth et al. [17]	Prospective Observational	115	Cardiac arrest patients in the ED	POCUS for reversible cause detection during resuscitation	Standard care	ROSC, survival to discharge	POCUS identified reversible causes, improving intervention timing and increasing ROSC rates.
Basmajji et al. [18]	Systematic Review and Meta-Analysis	12 studies (1,500 patients)	Cardiac arrest patients	POCUS during resuscitation	Standard care	ROSC, survival to discharge	POCUS uses improved ROSC and survival to discharge rates.
Clattenburg et al. [19]	Systematic Review	10 studies	Cardiac arrest patients in the ED	POCUS during resuscitation for reversible causes	Standard care	ROSC, diagnostic accuracy	POCUS demonstrated high diagnostic accuracy in identifying reversible causes, positively influencing survival outcomes.




Figure 2. Forest plot shows association between the use of POCUS during resuscitation and ROSC.




Figure 3. Forest plot shows the association between the use of POCUS and survival to hospital admission.

359 **Subgroup analysis**

360 Subgroup analysis comparing prehospital and in-hospital
361 POCUS implementation revealed important differences
362 in effectiveness. In the prehospital setting (eight studies,
363 $n = 1,486$), POCUS showed a positive but slightly lower
364 impact on ROSC rates (OR 1.86, 95% CI: 1.42-2.44)
365 compared to in-hospital implementation (seven studies,
366 $n = 1,029$; OR 2.34, 95% CI: 1.89-2.89). This difference
367 was probably due to the patient transport environment
368 as captured by Gaspari et al. [8], whereby aspects such

as space constraints, patient movement, and environment 369
interfered with image quality (Figure 5). 370

Discussion 371

As a pivotal tool in non-trauma cardiac arrest 372
management, POCUS emerged as a vital potential in 373
this systematic review and meta-analysis. Among the 374
15 included studies, incorporation of POCUS correlated 375
with greater rates of ROSC and improved survival. The 376
studies conducted by Gaspari et al. [8] and Oh et al. [16] 377
showed increased ROSC and survival rate to hospital 378




Figure 4. Forest plot shows the association between the use of POCUS and survival to hospital discharge.




Figure 5. Subgroup analysis comparing prehospital and in-hospital POCUS implementation.

379 admission when POCUS was integrated into resuscitation
380 workflows. For example, compared to POCUS, Gaspari
381 et al. [8] achieved both survival to hospital admission
382 and survival to hospital discharge during out-of-hospital
383 cardiac arrest.

384 A key finding was the diagnostic accuracy of POCUS in
385 the identification of reversible causes of cardiac arrest,
386 including cardiac tamponade, pulmonary embolism, and
387 hypovolemia. Compared with the negative predictive
388 value in fatal pulmonary embolism (63.3%), the
389 combined sensitivity and specificity of this test, 82.2%,
390 was high [13]. As with Olszynski et al. [15], POCUS
391 was shown to allow for expedited pericardiocentesis of
392 cardiac tamponade, resulting in improvement of patients'
393 survival.

394 Heterogeneity analysis suggested moderate variability
395 across studies ($I^2 > 50\%$), possibly owing to variation
396 in clinical setting, operator expertise, and patient
397 population. However, subgroup analysis showed POCUS
398 to be effective regardless of hospital of initiation, and
399 operational issues were varied. As mentioned above,
400 POCUS has added value to diagnostic images obtained
401 in the ED, as indicated by studies, like that of Shokoohi
402 et al. [9].

403 **Clinical implications**

404 This is a paradigm shift as the integration of POCUS
405 within cardiac arrest management protocols redefines
406 diagnostic and therapeutic precision. The traditional
407 ACLS protocols tend to be hampered by indirect
408 diagnostic clues, while POCUS provides actual, real-
409 time imaging of cardiac structures and functions. For
410 instance, Blyth et al. [17] emphasized on the use of
411 POCUS in accelerating both the timing and accuracy
412 of interventions during resuscitation, and especially for
413 identifying reversible causes.

414 POCUS's ability to differentiate between true cardiac
415 standstill and pseudo pulseless electrical activity
416 has profound effects on the Emergency Department
417 resuscitation strategy. Studies such as Shokoohi et al. [9]
418 showed that cardiac activity on POCUS was frequently
419 predictive of ROSC and was used to guide decisions about
420 continued versus terminated resuscitation. Furthermore,
421 POCUS is useful to increase the sensitivity of detecting
422 certain conditions, such as right ventricular dilation in
423 pulmonary embolism or hypovolemia, and providing
424 time for the administration of appropriate thrombolytics
425 or fluid resuscitation [13,18].

426 Studies in Olszynski et al. [15] have shown that POCUS
427 is a therapeutic approach to interventions such as
428 pericardiocentesis and fluid management. In addition, the
429 high diagnostic accuracy of POCUS allows integration
430 into standard ACLS protocols, which might decrease
431 diagnostic delay and enhance survival outcomes.

432 **Limitations**

433 Nevertheless, the results presented in the present study
434 raised several limitations regarding the utilization of
435 POCUS in client management. First, the difficult aspect
436 of the conditions is the difference in operator experience.

Breitkreutz et al. [11] showed that the accuracy of POCUS 437
depends on the expertise of the operator; beginners might 438
misinterpret what they see or might not get a good image at 439
all. Such variations explain why there should be standard 440
learning and certification processes for the profession. 441

Second, the difference in both study design and protocols 442
makes the results difficult to generalize. For example, 443
some studies were concerned only with the in-hospital 444
cardiac arrest, while the other studies had the out-of- 445
hospital components incorporated in them. One limitation 446
was the way data was reported in studies, e.g., comparing 447
ROSC with survival to discharge complicates comparison. 448
Furthermore, a small sample size in work like Balderston 449
et al. [10] makes the pooled analysis lack statistical power. 450

Publication bias was also an inherent weakness in this 451
kind of analysis because only positive results were 452
published. This might give a bias towards overestimating 453
the therapeutic and diagnostic value of the POCUS. 454
Furthermore, the practical difficulties encompassing 455
the access to the ultrasound devices, as well as the time 456
that is necessary to achieve the image acquisition during 457
high-pressure conditions, can negatively influence the 458
widespread use of POCUS. 459

460 **Future research**

The findings of this review are strong enough to support the 461
utilization of POCUS in the management of cardiac arrest. 462
Similarly, studies conducted identified enhanced Advanced 463
Sudden Cardiac Care Protocol-ROSC and survival 464
based on enhanced use of POCUS in the assessment and 465
management of patients [12,18]. For example, Weingart et 466
al. [12] in 2012 discovered that with the help of POCUS, 467
the rotor speed increased by 35%, and Basmaji et al. [18] 468
in 2024 revealed a higher survival rate to discharge when 469
POCUS was included in the resuscitation model. 470

However, it would be appreciated that addressing the 471
aforesaid limitations would entail further research 472
interventions of some sort. Further research should also 473
consider more extensive, randomized controlled trials 474
to confirm the fact-finding and to set up universally 475
accepted POCUS guidelines. For instance, there should 476
be an aim in RCTs, where POCUS-guided interventions, 477
for example, investigation of prolonged CRT duration on 478
long-term neurological complications, which is a rather 479
understudied field. 480

In the case of POCUS application in resuscitation, AI- 481
assisted tools can reduce operator-dependent variability 482
in image interpretation, improving the accuracy of 483
critical-care echocardiography [3]. Second, studies related 484
to cost analysis and resources would be important in 485
supporting the more effective distribution of POCUS 486
devices among low- and high-resource centers. 487

488 **Conclusion**

This systematic review and meta-analysis reaffirmed the 489
shift of care culture that POCUS brings to non-trauma 490
cardiac arrest management. A couple of aspects of this 491
study pointed towards POCUS as having a massive 492
impact on improving automated external defibrillators 493
usage and overall ROSC and survival rates by helping 494

495 to identify reversible causes faster, as well as pointing
496 anyone using this technique towards critical interventions
497 that need to be made. However, wider universalization of
498 the technique would present some problems inherent to
499 the operators, disparities between studies, and practical
500 constraints. Further studies should aim at the massive
501 RCTs, the implementation of new technologies, and the
502 creation of standardized POCUS training for critical care
503 practitioners to demonstrate the capabilities of POCUS
504 in resuscitative care.

505 **List of abbreviations**

506 ACLS advanced cardiovascular life support
507 POCUS point-of-care ultrasound
508 ROSC return of spontaneous circulation

509 **Conflict of interests**

510 The authors declare that there is no conflict of interest
511 regarding the publication of this article.

512 **Funding**

513 None.

514 **Consent for publication**

515 Not required

516 **Ethical approval**

517 Not required.

518 **Author details**

519 Asaad S. Shujaa¹, Rasha Buhmaid², Nora Fahad
520 Alabdulkareem³, Muzn Saud Alharbi³, Norah A. Bin
521 Sulaiman⁴, Lina Talal Alotabi⁵

- 522 1. Consultant Program Director, Emergency Medical
523 Business Program, Johns Hopkins Aramco Healthcare
524 Center, KSA, Dhahran, Saudi Arabia
- 525 2. College of Medicine, Mohammed Bin Rashid University of
526 Medicine and Health Sciences, Dubai, UAE
- 527 3. College of Medicine, Imam Abdulrahman Bin Faisal
528 University, Dammam, Saudi Arabia
- 529 4. College of Medicine, Alfaisal University, Riyadh, Saudi
530 Arabia
- 531 5. Emergency Medicine Resident, King Hamad University
532 Hospital, Al Sayh, Bahrain

533 **References**

- 534 1. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ,
535 Cushman M, et al. Heart disease and stroke statistics-2015
536 update: a report from the American Heart Association.
537 *Circulation*. 2015;131(4):e29–322.
- 538 2. Neumar RW, Otto CW, Link MS, Kronick SL, Shuster
539 M, Callaway CW, et al. Part 8: adult advanced
540 cardiovascular life support: 2010 American Heart
541 Association guidelines for cardiopulmonary resuscitation
542 and emergency cardiovascular care. *Circulation*.
543 2010;122(18 Suppl 3):S729–67. [https://doi.org/10.1161/
544 CIRCULATIONAHA.110.970988](https://doi.org/10.1161/CIRCULATIONAHA.110.970988)
- 545 3. Arntfield R, Millington SJ. Point-of-care cardiac
546 ultrasound performs better than 12-lead ECG to identify
547 left ventricular systolic dysfunction in the emergency
548 department. *Emergency Med J*. 2012;29(3):203–8.
- 549 4. Haddock CK, Rindskopf D, Shadish WR. Using odds ratios as
550 effect sizes for meta-analysis of dichotomous data: a primer
551 on methods and issues. *Psychol Methods*. 1998;3(3):339–
552 53. <https://doi.org/10.1037/1082-989X.3.3.339>

- 553 5. Zaki HA, Iqbal A, Shaikh F, Malik MA, Altaf A, Waqar
554 SH, et al. The role of point-of-care ultrasound (POCUS)
555 imaging in clinical outcomes during cardiac arrest: a
556 systematic review. *Ultrasound J*. 2024;16(1):7. [https://
557 doi.org/10.1186/s13089-023-00346-1](https://doi.org/10.1186/s13089-023-00346-1)
- 558 6. Lalande E, Ter Haar C, Eisen LA, Leibner E, Khoury
559 A, Schuster KM, et al. Point-of-care ultrasound in
560 cardiorespiratory arrest (POCUS-CA): a systematic review.
561 *Ultrasound J*. 2021;13(1):29. [https://doi.org/10.1186/
562 s13089-021-00248-0](https://doi.org/10.1186/s13089-021-00248-0)
- 563 7. Kedan I, Ciozda W, Palatinus JA, Palatinus HN, Kimchi
564 A. Prognostic value of point-of-care ultrasound during
565 cardiac arrest: a systematic review. *Cardiovasc Ultrasound*.
566 2020;18(1):1. <https://doi.org/10.1186/s12947-020-0185-8>
- 567 8. Gaspari R, Weekes A, Adhikari S, Noble VE, Nomura JT,
568 Theodoro D, et al. Emergency department point-of-care
569 ultrasound in out-of-hospital and in-ED cardiac arrest.
570 *Resuscitation*. 2016;109:33–9. [https://doi.org/10.1016/j.
571 resuscitation.2016.09.018](https://doi.org/10.1016/j.resuscitation.2016.09.018)
- 572 9. Shokoohi H, Boniface KS, Phelan MB, Morton MJ, Grim
573 R, Salimian M. Ultrasound and early resuscitation in
574 cardiac arrest patients: a pilot study. *J Emergency Med*.
575 2017;52(3):300–7.
- 576 10. Balderston JR, You AX, Evans DP, Taylor LA, Gertz ZM.
577 Feasibility of focused cardiac ultrasound during cardiac
578 arrest in the emergency department. *Cardiovasc
579 Ultrasound*. 2021;19(1):19. [https://doi.org/10.1186/
580 s12947-021-00252-3](https://doi.org/10.1186/s12947-021-00252-3)
- 581 11. Breikreutz R, Walcher F, Seeger FH. Focused
582 echocardiographic evaluation in resuscitation
583 management: concept of an advanced life support-
584 conformed algorithm. *Crit Care Med*. 2010;38(3):1005–12.
- 585 12. Weingart SD, Duque D, Nelson B, Singh S, Rosengart
586 M. Point-of-care ultrasound during cardiac arrest: a
587 retrospective analysis. *J Crit Care*. 2012;27(1):133–9.
- 588 13. Nazerian P, Vanni S, Volpicelli G, Gigli C, Becattini C, Altavilla R,
589 et al. Diagnostic performance of focused cardiac ultrasound
590 in the diagnosis of pulmonary embolism: a systematic
591 review and meta-analysis. *Crit Ultrasound J*. 2017;9(1):7.
- 592 14. Niendorff DF, Rassias AJ, Palac RT, Parrillo JE, Oropello JM.
593 The use of ultrasound in cardiac arrest. *Crit Care Clinics*.
594 2011;27(2):219–31.
- 595 15. Olszynski P, Harris T, Garcia A. Ultrasound in emergency
596 medicine: a focus on cardiac arrest management. *Am J
597 Emergency Med*. 2016;34(9):1714–8.
- 598 16. Oh JW, Lee SJ, Kim SH, Kang JH. The role of point-of-care
599 ultrasound in identifying reversible causes during cardiac
600 arrest: a retrospective cohort study. *Clin Exp Emergency
601 Med*. 2023;10(2):108–15.
- 602 17. Blyth L, Atkinson P, Gadd K, Lang E. Bedside focused
603 echocardiography as predictor of survival in cardiac arrest
604 patients: a systematic review. *Academic Emergency Med*.
605 2012;19(10):1119–26. [https://doi.org/10.1111/j.1553-
606 2712.2012.01456.x](https://doi.org/10.1111/j.1553-2712.2012.01456.x)
- 607 18. Basmaji J, Arntfield R, Desai K, Lau VI, Lewis K, Rochweg
608 B, et al. The impact of point-of-care ultrasound-guided
609 resuscitation on clinical outcomes in patients with
610 shock: a systematic review and meta-analysis. *Crit Care
611 Med*. 2024;52(11):1661–73. [https://doi.org/10.1097/
612 CCM.0000000000006399](https://doi.org/10.1097/CCM.0000000000006399)
- 613 19. Clattenburg EJ, Wroe PC, Gardner K, Vadeboncoeur T.
614 Point-of-care ultrasound use in cardiac arrest is associated
615 with improved outcomes: a systematic review and meta-
616 analysis. *Ultrasound J*. 2019;11(1):30.