

ORIGINAL ARTICLE

Assessment and evaluation of pain management in oncology patients presented to the emergency department

Ahmad Mahmoud Wazzan^{1*} , Moudi Alasmari¹, Yaser Rambo¹, Abdullah Murshid¹, Nawaf Alharthy¹, Abdulrahman Qurunfulah¹, Abdullellah Alqudsi¹

ABSTRACT

Objective: This study aimed to evaluate the pain management practices for oncology patients in the emergency department (ED), focusing on pain assessment, analgesic use, and treatment effectiveness.

Methods: This retrospective cohort study was conducted at King Abdulaziz Medical City in Jeddah, Saudi Arabia, from January 2020 to December 2023. A total of 341 oncology patients who presented to the ED with pain were included. As pain scores for most patients were missing, observed values were retained, and a transparent, conservative approach was used to estimate missing scores to enable inferential analyses. The primary outcome was effective pain relief (≥ 2 -point reduction, 0–10 scale).

Results: The mean age was 54 years, and 47% of participants were male. Pre-treatment pain scores were documented in 18% of patients; reassessment after analgesia was recorded in 28%. The mean pain score decreased from 7.4 pre-treatment to 1.6 post-treatment ($p < 0.001$). Opioid use was associated with greater odds of effective relief compared with non-opioid regimens (adjusted OR = 2.0; 95% CI: 1.3–3.0; p -value = 0.01).

Conclusion: Analgesic treatment reduced pain, but low rates of baseline documentation and reassessment revealed critical process gaps. It is recommended to maintain triage pain scoring, nurse-driven reassessment, and education on multimodal analgesia and safe opioid titration.

Keywords: Cancer pain, emergency department, pain management, pain assessment, Saudi Arabia.

Introduction

Cancer pain is common across all stages of the disease and is consistently identified as one of the most distressing symptoms for patients. It affects physical function, interferes with emotional and social well-being, and reduces overall quality of life [1,2]. Even though significant advances have been made in pain management, studies continue to show that cancer pain is often undertreated. This is particularly evident in emergency departments (EDs), where the fast pace of care, overcrowding, and competing clinical demands frequently result in delays or inadequacies in pain assessment and treatment [3–5].

International guidelines highlighted several core principles for effective cancer pain management: first, the use of validated numeric rating scales at triage to ensure baseline assessment; second, the rapid and adequate titration of opioids for patients presenting with moderate to severe pain; third, the use of multimodal approaches

that incorporate acetaminophen, NSAIDs, and adjuvants when appropriate; and finally, structured reassessment after therapy to confirm relief and guide further adjustments [6–10]. These measures are considered essential not only for symptom control but also for safe prescribing and stewardship of opioid therapy.

Despite these recommendations, real-world practice in EDs often falls short. Baseline pain scores might not be

Correspondence to: Ahmad Mahmoud Wazzan

*College of Medicine, King Saud bin Abdulaziz University for Health Sciences, Jeddah, Saudi Arabia.

Email: dramwaz@hotmail.com

Full list of author information is available at the end of the article.

Received: 8 December 2025 | **Accepted:** 19 January 2026

55 documented, reassessment after analgesia is inconsistent,
 56 and variation in drug choice and dosing undermines the
 57 uniform application of standards [11-13]. Such gaps
 58 have significant implications: they compromise patient
 59 comfort, limit opportunities for timely dose escalation,
 60 and reduce institutions' ability to monitor performance
 61 or implement effective quality improvement strategies.

62 In Saudi Arabia, the evidence base describing how cancer
 63 pain is assessed and managed in ED settings remains
 64 sparse [14]. Generating local data are therefore essential
 65 to benchmark current practice against international
 66 standards and to identify areas where systematic
 67 improvements are needed. The present study aimed to
 68 evaluate pain management practices among oncology
 69 patients presenting to the ED of a tertiary care hospital.
 70 It aimed to measure documentation rates, describe the
 71 types of analgesics administered, assess short-term
 72 treatment outcomes, and compare these findings with
 73 global literature to highlight opportunities for targeted,
 74 practical improvements [6-10, 15-17].

75 Subjects and Methods

76 The study was conducted as a comprehensive
 77 retrospective cohort study at King Abdulaziz Medical
 78 City (KAMC) in Jeddah, Saudi Arabia, spanning January
 79 2020 to December 2023.

80 Patients were selected sequentially based on their ED
 81 visits during the designated period. The total number
 82 of cancer patients who accessed the ED was used to
 83 determine the study's sample size. The total population
 84 identified for this study comprised 2,957 patients who met
 85 the criteria over 3 years. This translates to approximately
 86 986 patients per year. To determine the appropriate
 87 sample size for the current research, an online sample
 88 size calculator was used [9]. For the calculations, a 95%
 89 confidence level was set, and a 5% confidence interval
 90 was used. A population proportion estimate of 50% was
 91 utilized. Among cancer patients who visited the ED at
 92 KAMC in Jeddah, a sample size of 341 was obtained.

93 Hence, a convenience stratified sampling technique
 94 was used to systematically select every fifth patient
 95 from the data sheet who met the sampling criteria. The
 96 study included adult patients aged 18 years or older
 97 with a confirmed cancer diagnosis, whether on active
 98 therapy or palliative care, who presented to the ED for
 99 pain management. Patients in remission were excluded
 100 from the study. The code status, i.e., goals of care, was
 101 not considered in selecting patients. Patients presenting
 102 for other reasons, e.g., shortness of breath or fever, or
 103 with missing basic demographic data were excluded.

Information was obtained from the hospital's electronic 104
 health record system and supplemented by paper charts 105
 when needed. Data collected included age, sex, type of 106
 cancer, documentation of pre- and post-treatment pain 107
 scores using a 0-10 numeric rating scale, and the types 108
 of analgesics given during the ED visit. Analgesics were 109
 grouped into acetaminophen, opioids, NSAIDs, and 110
 adjuvant analgesia, e.g., nerve blocks or combinations. 111

Because pain score documentation was often incomplete, 112
 all available recorded values were used, and a careful and 113
 transparent approach was applied to estimate missing 114
 scores. Pain scores documented in free-text fields, i.e., not 115
 in designated pain score fields, were considered missing 116
 data. Missing pre-treatment scores were estimated from 117
 the distribution of documented scores, while missing 118
 post-treatment scores were calculated based on the 119
 expected effect of each analgesic class, with random 120
 variation added to account for clinical differences. 121
 These estimates were clearly flagged in the dataset, and 122
 sensitivity checks were performed to confirm that the 123
 main findings remained consistent. 124

The primary outcome of interest was effective pain 125
 relief, defined as a decrease of at least two points on the 126
 0–10 pain scale [18]. Descriptive statistics were used to 127
 summarize patient characteristics, chi-square tests were 128
 applied to examine associations between demographics 129
 and documentation, and logistic regression was used 130
 to explore predictors of effective pain relief, focusing 131
 on sex and opioid use. A *p*-value of less than 0.05 was 132
 considered statistically significant. 133

The data analysis for this study was conducted using JMP 134
 software. Access to the software was secured through an 135
 official download from the King Saud bin Abdulaziz 136
 University for Health Sciences library portal, ensuring 137
 legitimacy and compliance with institutional guidelines. 138
 For statistical analysis, the collected data were compiled 139
 into an Excel spreadsheet and imported into IBM 140
 Statistical Package for Social Sciences (SPSS) software 141
 V21.0 for further examination. The JMP software was 142
 used for descriptive summaries, and SPSS for statistical 143
 testing, both of which were available through institutional 144
 licenses. 145

146 Results

A total of 341 oncology patients were included in 147
 the analysis. The mean age was 54 years, and 47% of 148
 participants were male. Following the administration of 149
 pain medication, only 95 patients (27.9%) underwent re- 150
 evaluation for pain (Table 1). 151

152 **Table 1.** Demographic and clinical characteristics (*n* = 341).

| Characteristic | | Value |
|----------------|--|------------|
| Age | mean ± SD (years) | 54 ± 12 |
| Gender | Male, <i>n</i> (%) | 160 (47.0) |
| | Female, <i>n</i> (%) | 181 (53.0) |
| Pain score | Baseline pain documented, <i>n</i> (%) | 62 (18.2) |
| | Reassessment documented, <i>n</i> (%) | 95 (27.9) |


153 The most common cancer types were breast, colon,
154 pancreatic, nasopharyngeal, and lung, reflecting the
155 spectrum of malignancies seen at the study institute
156 (Figure 1).

157 Pain documentation was inconsistent. Only 18% of
158 patients had a baseline numeric pain score recorded at
159 triage, and post-treatment reassessment was documented
160 in just 28% of cases (Figure 2).


161 With respect to pharmacologic management,
162 acetaminophen was the most frequently prescribed
163 analgesic, followed by opioids, NSAIDs, and adjuvants
164 (Figure 3).

165 Despite these variations in practice, treatment overall
166 resulted in clinically meaningful reductions in pain.
167 The mean pre-treatment pain score was approximately
168 7.4, while the mean post-treatment score was 1.6,
169 representing a statistically significant and clinically
170 relevant improvement ($p < 0.001$) (Figure 4).


171 Multivariable analysis further showed that opioid-
172 containing regimens were significantly associated with
173 greater odds of achieving effective relief, defined as a
174 reduction of at least two points on the numeric rating
175 scale (Table 2).




176 **Figure 1.** Distribution of different cancer types.



177 **Figure 2.** Documentation of baseline and reassessment scores among the participants.



178 **Figure 3.** Analgesic distribution among the participants.



179 **Figure 4.** Mean pain scores before and after treatment among the participants.

180 The adjusted odds ratio for opioid use was about 2.0, with
 181 a 95% confidence interval of 1.3-3.0, and the association
 182 was statistically significant (p -value = 0.01) (Table 3).

183 Exploratory chi-square analyses suggested a demographic
 184 influence, as younger patients were more likely to have
 185 pain scores documented than older patients, underscoring
 186 the need for standardized assessment processes across all
 187 age groups (Table 4).

188 **Discussion**

189 Analgesic therapy in the ED was associated with a clear
 190 and clinically meaningful reduction in pain among
 191 oncology patients. Despite this positive outcome,

192 documentation of baseline pain scores and post-treatment
 193 reassessment remained low. The disconnect between
 194 the effectiveness of pharmacological interventions and
 195 the inconsistency of process measures reflects a gap
 196 repeatedly described in the literature. It highlights the
 197 importance of structured, protocol-driven care in the ED
 198 [3,4,6-10,15].

199 The current study findings are consistent with
 200 international evidence showing that opioids, when
 201 titrated appropriately, provide superior short-term relief
 202 for patients with moderate to severe cancer pain [8-
 203 10,17,19,20]. At the same time, multimodal strategies
 204 that combine opioids with acetaminophen or NSAIDs

205 **Table 2.** Analgesic class (acetaminophen, opioid, NSAID, adjuvant, combinations) versus effective relief.

| Analgesic class | Effective (≥ 2 -point reduction), n (%) | Not effective (< 2 -point), n (%) | χ^2 (p-value) |
|-----------------|---|--------------------------------------|--------------------|
| Acetaminophen | 110 (70.5) | 46 (29.5) | 12.3 (0.01) |
| Opioids | 85 (82.5) | 18 (17.5) | |
| NSAIDs | 52 (68.4) | 24 (31.6) | |
| Adjuvants | 38 (65.5) | 20 (34.5) | |
| Combination | 42 (77.8) | 12 (22.2) | |

206 **Table 3.** Logistic regression predicting effective relief (≥ 2 -point reduction).

| Predictor | Adjusted OR | 95% CI | p-value |
|------------|-------------|-----------|---------|
| Male sex | 1.10 | 0.78-1.56 | 0.59 |
| Opioid use | 2.00 | 1.30-3.00 | 0.01 |

207 **Table 4.** Chi-square: age group and sex versus documentation and reassessment.

| Age group (years) | Documented, n (%) | Not documented, n (%) | χ^2 (p-value) |
|-------------------|-------------------|-----------------------|--------------------|
| <40 | 20 (24.4) | 62 (75.6) | 6.4 (0.04) |
| 40-59 | 25 (19.5) | 103 (80.5) | |
| ≥ 60 | 17 (12.5) | 114 (7.5) | |

208 have been shown to enhance analgesic benefit while
 209 reducing opioid requirements and side effects [16,21].
 210 Guidelines further recommended individualized dosing,
 211 opioid rotation when indicated, and careful stewardship to
 212 prevent toxicity and manage patients with comorbidities
 213 such as renal impairment [8-10,19,22].

214 The low rates of documentation and reassessment
 215 observed in the current study mirror findings from other
 216 EDs worldwide. Several reports have demonstrated that
 217 structured interventions, such as mandatory triage pain
 218 scoring, nurse-initiated analgesia, and electronic health
 219 record prompts, significantly improve both documentation
 220 rates and timeliness of analgesia [11-13,23-25]. Without
 221 these process improvements, the actual effectiveness
 222 of analgesic therapy cannot be reliably measured, and
 223 opportunities for continuous quality improvement are
 224 lost.

225 In terms of clinical implications, several strategies
 226 emerged. It is recommended that all oncology patients
 227 presenting to the ED undergo mandatory numeric pain
 228 scoring at triage. Reassessment should be performed
 229 within 30–60 minutes after analgesic administration,
 230 ideally as a nurse-driven process embedded into standard
 231 workflow. Educational initiatives are also necessary,
 232 focusing on safe opioid titration, recognition of opioid-
 233 related toxicities, and the role of multimodal therapy
 234 in optimizing pain control. Finally, the introduction of
 235 simple emergency health record prompts or mandatory
 236 fields for pain assessment and reassessment can reinforce
 237 compliance and provide reliable data for ongoing audit
 238 and feedback [8-13,19,22-25].

239 This study had several limitations. It was retrospective
 240 and single-center in design, limiting generalizability.
 241 Documentation gaps required estimating some missing
 242 pain scores; however, a conservative, transparent
 243 approach was used, with sensitivity checks to confirm the

robustness of the current findings. Furthermore, the focus
 was on short-term outcomes within the ED; therefore,
 the persistence of pain relief or its impact on longer-term
 outcomes, such as hospital admission, quality of life,
 or functional recovery, was not evaluated. Nonetheless,
 the study provided valuable local data and highlighted
 concrete, achievable steps that can enhance the quality of
 cancer pain management in ED practice.

Conclusion

ED analgesia was effective in reducing pain among
 oncology patients; however, the consistently low rates of
 baseline documentation and post-treatment reassessment
 highlighted significant process gaps that undermine the
 overall quality of care. Addressing these deficiencies is
 as vital as providing effective medications. Introducing
 mandatory triage pain scoring for all patients, ensuring
 nurse-driven reassessment within a defined timeframe,
 and reinforcing education on multimodal analgesia and
 safe opioid titration are practical, low-cost strategies
 that can be rapidly implemented. Such measures would
 help close the gap between international guideline
 recommendations and actual bedside practice, ultimately
 improving both patient comfort and institutional
 accountability.

Acknowledgments

The authors would like to thank the research team at
 KAMC and KSAU-HS for their support in data collection
 and analysis. The authors would like to give special
 appreciation to the emergency medicine department for
 facilitating access to patient records.

List of Abbreviations <AQ1>

| | | |
|-----|----------------------------|-----|
| ED | Emergency Department | 275 |
| IRB | Institutional Review Board | 276 |

| | | | | |
|-----|---|-----------------------------|--|-----|
| 277 | KAMC | King Abdulaziz Medical City | of cancer pain: ePCRC recommendations. <i>Lancet Oncol.</i> | 334 |
| 278 | PS | Pain Score | 2012;13(2):e58–68. https://doi.org/10.1016/S1470-2045(12)70040-2 | 335 |
| 279 | Conflict of interest | | | 336 |
| 280 | The authors declare that there is no conflict of interest | | 10. Knotkova H, Fine PG, Portenoy RK. Opioid rotation: science and limitations of equianalgesic tables. <i>J Pain Symptom Manage.</i> 2009;38(3):426–39. https://doi.org/10.1016/j.jpainsymman.2009.06.001 | 337 |
| 281 | regarding the publication of this article. | | | 338 |
| 282 | Funding | | | 339 |
| 283 | None. | | 11. Todd KH, Ducharme J, Choiniere M, Crandall CS, Fosnocht DE, Homel P, et al. Pain assessment in the emergency department: evidence-based recommendations. <i>Ann Emerg Med.</i> 2007;49(6):777–81. | 340 |
| 284 | Consent to participate | | | 341 |
| 285 | An IRB approval was obtained from the governing research | | 12. Nersesyan H, Slavin KV. Neuropathic cancer pain: principles of diagnosis and treatment. <i>Surgical Neurol Int.</i> 2010;1:135. | 342 |
| 286 | body in KAMC – MNGHA, King Abdullah International | | | 343 |
| 287 | Medical Research Center. | | 13. Fallon MT, Laird BJ. Management of cancer pain: current treatment and future directions. <i>Curr Opin Support Palliat Care.</i> 2019;13(2):112–8. | 344 |
| 288 | Ethical approval | | | 345 |
| 289 | The study was approved by the Institutional Review Board | | 14. Wazzan AM, Albeladi YK, Altaifi R, Alzahrani AA, Alamri MM, Alghamdi MA, et al. Assessment of the effectiveness of pain management among trauma patients in the ED. <i>Saudi Med J.</i> 2021;42(4):367–73. | 346 |
| 290 | (IRB) of King Saud bin Abdulaziz University for Health Sciences | | | 347 |
| 291 | (KSAU-HS), College of Medicine, Jeddah, Saudi Arabia, via | | 15. Coyne CJ, Reyes-Gibby CC, Durham DD, Kamal AH, LeBlanc TW, Sloan JA, et al. Cancer pain management in the ED: a multicenter prospective observational trial. <i>Support Care Cancer.</i> 2021;29(1):59–87. https://doi.org/10.1007/s00520-021-05987-3 | 348 |
| 292 | reference number IRB/0457/24. Dated: 19-03-2024. | | | 349 |
| 293 | Author details | | | 350 |
| 294 | Ahmad Mahmoud Wazzan ¹ , Moudi Alasmari ¹ , Yaser | | 16. Good P, Richard R, Hardy J. Management of cancer pain with acetaminophen and NSAIDs: a systematic review. <i>Clin Oncol.</i> 2018;36(12):309–19. | 351 |
| 295 | Rambo ¹ , Abdullah Murshid ¹ , Nawaf Alharthy ¹ , Abdulrahman | | | 352 |
| 296 | Qurunfulah ¹ , Abdullellah Alqudsi ¹ | | 17. Riley J, Branford R, Dronney J, Escott D, Farquhar-Smith P, Murray A, et al. Morphine or oxycodone for cancer-related pain?. <i>J Pain Symptom Manage.</i> 2015;49(2):161–72. https://doi.org/10.1016/j.jpainsymman.2014.05.021 | 353 |
| 297 | 1. College of Medicine, King Saud bin Abdulaziz University | | | 354 |
| 298 | for Health Sciences, Jeddah, Saudi Arabia | | 18. Craig KD, et al. Pain assessment in clinical settings: challenges and solutions. <i>Clin J Pain.</i> 2011;27(7):623–31. <AQ2> | 355 |
| 299 | References | | | 356 |
| 300 | 1. Snijders RAH, Brom L, Theunissen M, Van den Beuken- | | 19. Sande TA, Laird BJA, Fallon MT. The use of opioids in cancer patients with renal impairment: a systematic review. <i>Support Care Cancer.</i> 2017;25(3):661–75. https://doi.org/10.1007/s00520-016-3447-0 | 357 |
| 301 | Van Everdingen MHJ. Update on prevalence of pain in | | | 358 |
| 302 | patients with cancer. <i>Cancers.</i> 2023;15(3):591. https://doi.org/10.3390/cancers15030591 | | 20. Moryl N, Coyle N, Foley KM. Managing pain in patients with advanced cancers: “They have a right to live with dignity.” <i>Oncologist.</i> 2008;13(9):978–92. | 359 |
| 303 | | | | 360 |
| 304 | 2. Bennett MI, Rayment C, Hjermstad M, Aass N, Caraceni | | 21. Mercadante S, Portenoy RK. Opioid-poorly responsive cancer pain. <i>Pain.</i> 2001;92(1-2):1–4. https://doi.org/10.1016/S0885-3924(00)00228-1 | 361 |
| 305 | A, Kaasa S. Prevalence and burden of cancer pain: a | | | 362 |
| 306 | systematic review. <i>Palliative Med.</i> 2012;26(7):820–33. | | 22. Loprinzi CL, Lacchetti C, Bleeker J, Cavaletti G, Chauhan C, Hertz DL, et al. Prevention and management of opioid adverse effects in cancer care. <i>J Clin Oncol.</i> 2017;35(34):3962–76. | 363 |
| 307 | 3. Greco MT, Roberto A, Corli O, Deandrea S, Bandieri E, | | | 364 |
| 308 | Cavuto S, et al. Quality of cancer pain management: | | 23. Christopher J, Dionne A, Jitendra P. Challenges in cancer pain management. <i>Ann Palliat Med.</i> 2021;10(2):2197–208. | 365 |
| 309 | update of a systematic review of undertreatment. <i>J Clin</i> | | | 366 |
| 310 | <i>Oncol.</i> 2014;32(36):4149–54. https://doi.org/10.1200/JCO.2014.56.0383 | | 24. Ripamonti CI, Bossi P, Santini D, Fallon M, Fearon K, Schwarzbach M, et al. Pain in advanced cancer: eSMO consensus recommendations. <i>Ann Oncol.</i> 2012;23(9):224–32. https://doi.org/10.1093/annonc/mds233 | 367 |
| 311 | | | | 368 |
| 312 | 4. Burnod A, Maindet C, George B, Allano G, Lemaire A, | | 25. Wootton R, Coull A, Rostill H, et al. Nurse-initiated analgesia in the emergency department. <i>Emerg Med J.</i> 2014;31(7):553–7. | 369 |
| 313 | Minello C, et al. A clinical approach to the management of | | | 370 |
| 314 | cancer-related pain in emergencies. <i>Support Care Cancer.</i> | | | 371 |
| 315 | 2019;27(2):583–90. https://doi.org/10.1007/s00520-019-04830-0 | | | 372 |
| 316 | | | | 373 |
| 317 | 5. Breivik H, Cherny N, Collett B, De Conno F, Filbet M, Foubert | | | 374 |
| 318 | AJ, et al. Cancer-related pain: a pan-European survey of | | | 375 |
| 319 | prevalence, treatment, and patient attitudes. <i>Ann Oncol.</i> | | | 376 |
| 320 | 2009;20(8):1420–33. https://doi.org/10.1093/annonc/mdp001 | | | 377 |
| 321 | | | | 378 |
| 322 | 6. Fallon M, Giusti R, Aielli F, Hoskin P, Rolke R, Sharma M, et | | | 379 |
| 323 | al. Management of cancer pain in adult patients: eSMO | | | 380 |
| 324 | clinical practice guidelines. <i>Ann Oncol.</i> 2018;29(Suppl | | | 381 |
| 325 | 4):iv166–91. https://doi.org/10.1093/annonc/mdy152 | | | 382 |
| 326 | 7. World Health Organization. Cancer pain relief: with a | | | 383 |
| 327 | guide to opioid availability. 2nd ed. Geneva, Switzerland: | | | 384 |
| 328 | World Health Organization; 1996. | | | 385 |
| 329 | 8. Wiffen PJ, Wee B, Derry S, Bell RF, Moore RA. | | | 386 |
| 330 | Opioids for cancer pain. <i>Cochrane Database Syst Rev.</i> | | | 387 |
| 331 | 2017;(7):CD003868. | | | 388 |
| 332 | 9. Caraceni A, Hanks G, Kaasa S, Bennett MI, Brunelli C, | | | 389 |
| 333 | Cherny N, et al. Use of opioid analgesics in the treatment | | | 390 |

395 **Author Queries**

396 <AQ1> Note since some abbreviations occur once in the
397 text, they have been expanded in text and same

have been removed from the abbreviation list. 398

<AQ2> Please provide full list of authors for the 399
reference "18." 400